

Em 2001, uma diretiva do Conselho e Parlamento Europeu referente ao uso de advertências de saúde em embalagens de cigarros se tornou lei. No mínimo advertências de texto devem aparecer os rótulos; o uso de imagens é opcional. A Comissão Europeia (EC) produziu várias advertências ilustradas e de texto que os países-membro podem optar por usar.

Data de implementação e rodadas adicionais 2004

Cobertura da embalagem

Os estados-membro da União Européia devem exibir advertências de texto segundo os padrões abaixo:

EXIGÊNCIA MÍNIMA

	sem bordas			com bordas		
	total	frente	verso	total	frente	verso
Países unilingües	35%	30%	40%	48%	43%	53%
Países bilingües	39%	32%	45%	52%	45%	58%
Países trilingües	43%	35%	50%	56%	48%	63%

Freqüência de rotação e histórico

Há uma série opcional com 42 advertências ilustradas. A freqüência recomendada de rotação não é especificada.

Restrições sobre informações enganosas


Descritores enganosos, como “light” e “suave”, são proibidos de figurarem nas embalagens.

Notas especiais

Muitos países que não são membros da União Européia também adotaram as imagens de advertências da comunidade para uso em suas embalagens.

ADVERTÊNCIAS

2004



ADVERTÊNCIAS

2004



Smoking can damage the sperm and decrease fertility



Smoking can damage the sperm and decrease fertility



Smoke contains benzene, nitrosamines, formaldehyde and hydrogen cyanide

The risk of having a stroke is approximately 50% higher in smokers than in non-smokers

Smoking clogs the arteries and causes heart attacks and strokes



Smoking when pregnant harms your baby



Smokers die younger



Smoking can cause a slow and painful death



Get help to stop smoking: 0803 00 00 00



Smoking causes fatal lung cancer



Smoking causes fatal lung cancer



Smoking causes fatal lung cancer

Stopping smoking reduces the risk of fatal heart and lung diseases




Smoking may reduce the blood flow and causes impotence



Smoking may reduce the blood flow and causes impotence



Smoking may reduce the blood flow and causes impotence



Smoking when pregnant harms your baby

You can do it, we can help

Your doctor or your pharmacist can help you stop smoking



Smoking causes ageing of the skin



Stopping smoking reduces the risk of fatal heart and lung diseases



Smoking causes ageing of the skin

Lifetime smokers lose an average of 14 years of life

Smokers die younger



Smokers die younger

0803 00 00 00

Choose freedom, we'll help you

Get help to stop smoking

Smoking is a serious nicotine addiction, don't be afraid to ask for help

Your doctor or your pharmacist can help you stop smoking

ADVERTÊNCIAS

2004

